Cimetidine	No effect on plasma concentrations or LDL-C lowering efficacy of atorvastatin 1 in TG-lowering effect of atorvastatin from	
Dittiazem Hydrochloride	34% to 26%. Steady-state diffiagem increases the	
Ditazem Hydrochonos	exposure, based on AUC _{LASTs} , of a single dose of atorvastatin by approximately 50%.	
Antipyrine	Altorvastatin had no effect on the PK of antipyrine.	Antipyrine was used as a non-specifi model for drugs metabolized by the microsomal hepatic enzyme syster (cytochrome P-450 system). Interaction with other drugs metabolized via the sam cytochrome iscozymes are not expected.
Macrolide Antibiotics (azithromycin, clarithromycin, erythromycin)	In healthy adults, coadministration of atomastatin (10mg QD) and azilfromycin (500mg QD) did not significantly after the plasma concentrations of atomastatin.	See WARNINGS AND PRECAUTIONS Muscle Effects; DETAILED PHARMACOL OGY - Human Pharmacokinetics
Clarithromycin and erythromycin are both CYP3A4 inhibitors	† plasma concentration by ~40% with erythromycin (500mg OID) and ~80% with clarithromycin (500mg BID) when coadministered with atorvastatin (10mg QD)	
Protesse Inhibitors	1 plasma concentrations of allorvastatin when abrovastatin 10 mg OD is coadministered with refillaruir meeylate 1250 mg BID 1, AUC by 14% and 1 Cmsx by 122% 1, AUC by 14% and 1 Cmsx by 127% 1, AUC by 19, 9 fold and 1 Cmsx by 4.7 fold with adrovastatin 20mg OD and Lopinaviv 1, AUC by 30 fold and 1 Cmsx by 4.3 fold with abrovastatin 40mg OD and Ritonavir 400mg IS augularuir 400mg IS augularuir 400mg ISD 1	Concordant use of strong inhibitors of CVP 3A4 with abrovastatin about the use with caution. If such use must be institute lower zarting and maintenance doese carting and maintenance doese carting asked to the considered an patients should be considered an patients should be monitored closel for musculcokeletal effects
Cyclosporine	Concomitant administration of atomistatin 10mg and cyclosporine 5.2mg/kg/day resulted in a 7.7 fold increase in exposure to atomistatin.	Concomitant use should be used wit caution. In cases where co-administratio of atonivastatin with cyclosportine is necessary, the dose of atonivastati should not exceed 10mg. Se WARNINGS and PRECAUTIONS Muscle Effects; DETAILED PHARMA COLOGY - Human Pharmacolonetos
tracenazole	Concernitant administration of atorvastain 20-40mg and traconazole 200mg daily resulted in a 2.5-3.3-4old increase in atorvastatin AUC.	Concomitant use of strong inhibitors CYP 3A4 with atorvestatin should be used with caution. If such use must be instituted, lower starring and maintenand doses of atorvestatin should be considered and patients should be considered and patients should be monitored closely for musculoskelett effects (see DETAILED PHARMACOL GGY - Human Pharmacokinetics).
Efavirenz	AUC by 41% and Cmax by 1% with atorvastatin 10mg and Etavirenz 600mg daily.	
Rfampin	Co-administration.": † AUC by 30% and † Cmsx by 2.7 fold co-administered atorvastatin 40mg single dose and Pitampin 600mg daily. Separate administration. † in AUC by 80% and † Cmsx by 40% with atorvastatin 40mg single dose and	Due to the clual interaction mechanism or ritampin, simultaneous co-administration of atorvastatin with rifampin is recommended, as delayed administration or atorvastatin after administration or rifampin has been associated with significant reduction in atorvastatin

with abstraction of long region loss and Inflammo Office and Infla

Fusidic Acid

**Date given as AAM change represent a simple and between constructation and administration later (a. 1.445 in on draps).

**Date given in a Visit great present a full-time condition by districtation later (b. 7. in ordinary).

**Date given in Visit great given the properties of th

Firmup 1 propertionel resistance and combined Minisch Dysiglogenia. Noticing Testinal Controlled Dispetitioners and Properties of the Controlled Dispetition of the Controlled Dispetition

num dose is 80 mg/day.

Sociage of Anonsathar should be individualized according the baseline LDL-C, total-CHDL-C ratio and/or TG levels to achieve the numeroded desired lipid values at the lowest dose needed to achieve LDL-C desired level. Lipid levels should be monitored disaly and, If necessary, the dose of Anonsatatian adjusted based on desired lipid levels recommended by galdelines.

Servero Displosterials
In patients with server dysliptionials, including homozygous and heterozygous familial hypercholesterolemia and dysbetalloporotein
In patients with server dysliptionials, including homozygous and heterozygous familial hypercholesterolemia and dysbetalloporotein
Interactions, Muscle Ethoris, DRUG INTERACTIONIS,
Interactions, Muscle Ethoris, DRUG INTERACTIONIS,
Interactions, Muscle Ethoris, DRUG INTERACTIONIS

On INTERACT

Interactions, Nusuel Effects; (PRIJ 6) INTERACTIONS).

Heteropropus Finality hypercholestrections for Pediatric Palinates (10.17 years of ago)
in the population, the incommended dating dose of Abrovastian in 10 insplicy, the maximum recommended dose is Dimplisy (dose
in the population, the incommended palinates of Abrovastian in 10 insplicy, the maximum recommended dose is Dimplisy (dose
commended palin deserve) (are INDICATIONS AND CLINICAL USE and PHARMACCLORY, Clinical Bluebase). Adjustments should
be made at Intervals of 1 weeks a more.

Perventional of Cenderical Chaeses

Preventional of Cenderical Chaeses

and the prevention of the production of the primary prevention of mycardial infection used a dose of 10 ing abrovastian once daily, for a secondary prevention of the procedure of the production of the primary prevention of mycardial infection used as dose of 10 ing abrovastian once daily, for a secondary prevention of mycardial infection and a dose of 10 ing abrovastian once daily, for a secondary prevention of mycardial infection of the production o

Concomitant Therapy See DRUG INTERACTIONS.

see invarious on our procusul invito.

OVERDOSAGE
There is no specific treatment for abrovastatin over dosage. Should an overdose occur, the patient should be treated symptomatically and supportive measures instituted as are required. Due to extensive drug binding to plasma proteins, hemodalysis is not expected to algorithms and sonoversation desirance (See ADVERSE REACTIONS).

e, contact your regional Poison Control Centre immediately. ACTION AND CLINICAL PHARMACOLOGY

ACTION AND CLIBINAL PHANIMALALANA

ACTION AND CLIBINAL PHANIMALALANA

AND CLIBINAL PHANIMALANA

Abstraction research (LDL).

Abstraction rese

associated with decreased HDLO or increased LDLO.

Epidemiologic, clinical and experimental studies have established that high LDLO, low HDLO and high plasma TG promote human atherocoleropis and are risk flastors for developing cardiovascular disease. Some studies have also shown that the total (TG)+HDLO and the product of commany array disease, some studies have also shown that the total (TG)+HDLO and the product of commany array disease, some contrast, increased levels of HDLO are associated with decreased cardiovascular risk. Drug therapies that reduce levels of LDLO or decrease TG while simultaneously increasing HDLO have demonstrated reductions in rates of cardiovascular mortality and mortality.

optotellisprocrienmina, accruatation has been shown to reduce levels of total industrient (Ital-Ci, LDC-C, op and for IET), and for information of the control of the contr In patients with type II hyperlipidemia, atorvastatin improved endothelial dysfunction. Atorvastatin significantly improved flow-nendothelium-dependent dilatation induced by reactive hyperemia, as assessed by brachial ultrascund (p-0.01).

embridation dependent distation in Account by resident hyperensis, as assessed by locational distancies of Account by Performance Accounts of the Performance Accounts of

attractabilities is likely to be accorded in human milk.

Methodisme. Amonative in settinely methodistical to other and pare-hydroxyllated derivatives by cyto-home P-493 3-M (CYP 244) and to virious bette adultion products. In his inhibition of HMD-CA inductions by other and pare-hydroxyllated methodistics is expeniented to settine better adultion products. In his inhibition of HMD-CA inductions by other and pare-hydroxyllated methodistics is sequelected by the control of the contr

approximately 14 hours, but the half-life for inhibitory activity for HMD-CAA medicates as in 20 to 30 to one of the three contribution of looper designation of the contribution of looper designation of looper design

Condender Plasma concentration of attendants in revenue file (responsible) (20 Kight of each of contract with young and contract with young and the contract with the contract

CAUTION
Foods, Drugs, Devices and Cosmetics Act prohibits dispensing without prescription

AVAILABILITY: Atorvast-Natrapharm is available in dosage strengths of 10 mg, 20 mg, 40 mg and 80 mg Atorvastatin.

PACKAGING: Alu-alu blister pack x 10's (Box of 10's and 100's)

For suspected adverse drug reaction, report to the FDA; www.fda.gov.ph

Registration Number: 10mg (DR-XY40122), 20mg (DR-XY40121), 40mg (DR-XY40120), 80mg (DR-XY40119) Date of First Authorization: 10mg, 20mg, 40mg, 80mg (Debber 2011)

STORE AT ROOM TEMPERATURES NOT EXCEEDING 30°C

Manufactured by Apotex,Inc. 150 Signet Drive, Imported by/Packed for Natrapharm, Inc. The Patriot Building, SLEX, Sucat, Parañaque City

IMPORTANT: PLEASE READ

Identify Ide

ABOUT THIS MEDICATION: PLEASE READ

What the medication is used for: our detor has prescribed here polls to help fower your cholesterol or other fats in the blood (such as triglyceristies) and to prevent and/ourscular disease such as heart attacks. High levels of cholesterol and other fats can cause heart disease by ologging the blood sessies that feed blood and oursen to the heart.

vessels that feed blood and cargons to the heart.

Clutture 10.17 years and aim heteroognomis serial by specificient coloring and a coloring to the parents) and a besent form on a first or that factor disclinational disclination as a determined by your doctor, on the heart factor of the factor o

The properties of the state of The contraction can be a second to the contraction of the contraction can be contracted to the contract of the contraction contracted to the contraction contracted to the contracted to the contracted to the contraction contracted to the contracted to the contraction contracted to the contracted t

Abrocation also reduces the risk of heart attacks and shokes in people with multiple risk factors for coronary heart disease such as highly bodd pressure and dislates. When used by people who have suffered a heart attack in the past, atomistation reduces the risk of haring another heart attack.

Available from the attack.

Accessations in only alwance by precipions that evering a obcoor.

When it shaded not be affected in the property of the several property of the several property of the several property of the indicated ingredient is and what the important non medicinal regredient is and what the important non medicinal regredient is an indicated in the several propertient series.

- New section liver disease or unexplained horizonase in hier entrymes.
- Assist properties the enter-desiring.

What the medicinal ingredient is: atoryastatin calcium propylene olycol solvati

storwateria celatum progrieme glycos torines.

What the imported momentical intereferent aces:
Abovesation it basits contenes caucium acestus, oreocumentose sodium, sodium carbonate, microcystatine cellulose, magne
Abovesation italianis, contenes caucium acestus, oreocumentose sodium, sodium carbonate, microcystatine cellulose, magne
Abovesation italianis, contenes in contene

What doseage from it. comes in Tribette are evaluated in strength of the stren

observative projects exercise
 observative projects exercise
 observative projects exercise
 observative projects
 observative projects

is with most medicines, interaction with other drugs is possible. Tell your doctor or pharmacist if you are taking any other medication scholing prescription, non-prescription and natural health products. In particular, these drugs may interact with atorvistatin:

concisions (common-like mediones) cylologopine (BANDMM/LEE); permittrooti (JCPIDI) germittrooti (JCPIDI) permittrooti (JCPIDI) pick-lowering doses of naiori (notortio acids) synthomycini, distrinopini or acids artifungia algentis (seloconazole or litaconazole) prittromycini, distrinopini or acids artifungia algentis (seloconazole or litaconazole) reflazodore (SERZO/NEE) SIGNAMI/S BLAFFI (CRINIA/NII), nafinavir mesylate (VIRACEPTRI, rinnavir NIPA

work (SERZONES)

WRS SULFATE (CRIXIVANS), netinavir mesylate (VIRACEPTII), ribnavir (NORVIRIS), sequinavir mesylate acci (a, CRIXIVANS), netinavir mesylate acci (a, CRIXIVANS), netinavir (a, KALETRAIS)

ASE^(**), loperaviritionavir (a, KALETRAIS)

- grapherius (see - especially if implesting parasities of 2 it idens of grapherius) pare at encire
 PROPER USE OF THIS MEDICATION
 We often cannot see or feel the problems that high cholesterol causes until a lot of time has passed. That's why it is important to take
 these palls guit as prescribed. You jurc child and your doctor will be watching youlyour child's cholesterol levels to get them doesn to a
 see for range. Here are even important tips.

late largos, hire are some important figur.

Frequency of the property of the

Abrovatative was prescribed by syndrout child only, Don't give these gifts to enyone relate.
 Abrovatative was prescribed by syndrout child only the prescribed by the

inseed Does:

you'your child miss take a pill, take it as soon as possible. But lif it is almost time for the next dose, skip the missed dose and just take SIDE EFFECTS AND WHAT TO DO ABOUT THEM

BIDE EFFECTS AND WHAT TO DO ABOUT THEM

Man people do not have any poblems with side effects when taking this modeline. However, all medicines can cause unwanted side
effects. Check with your door or pharmacets promptly if any of the bollowing persist or become thousesome:

- conclusion/cultur-league
- conclusion/cultur-league
- inclusion-cultur-league
-- domain pain or uppert
- domain pain or uppert
- unaffing of throwing up
- working of throwing up
- where years, a few poperations of the late year.
- they range, a few poperations of the late year.
- they remain the process of the late year.
- the process best deference when once extractive parameter cough another shariness of breath or fever.
- treating proteines recluding parameter cough another shariness of breath or fever.
- the process of the p

Symptom/Effect		Talk with your doctor or pharmacist		Stop taking the drug and seek
		Only if severe	In all cases	emergency medical assistance
Rare	Muscle pain that you cannot explain		√	
	Muscle tenderness or weakness		√	
	Generalized weakness, especially if you don't feel well		V	
	Brownish or		- √	

HOW TO STORE IT

Always keep medicine well out of the reach of children.

Keep atorvastain at room temperature (15-90°C), away from

The control of the pattern of the pa

Email details to jssandlego@
 Fax details to (02)821-7383
 Or call us at (02)821-7382

MORE INFORMATION
For more information, please contact your doctor, pharmacist or other healthcare professional.

This leaflet plus the full product monograph, prepared for health professionals, can be obtained by contacting DISpedia, Applox's Drug Information Service at:

This leaflet was prepared by Apotex Inc., Toronto, Ontario, M9L 1t9 Date of Revision: April 27, 2011