

Natrapharm



# Ambroxol HCl

## Zobrixol<sup>®</sup>

75mg Sustained-Release Capsule

Mucolytics

### Description of the Product

Ambroxol HCl (Zobrixol) 75mg capsule is white to off-white to slightly yellowish pellets encapsulated empty gelatin capsule size #2 with brown opaque cap and transparent body.

### What is in the Medicine?

This medicine contains Ambroxol, which stimulates respiratory fluid secretion, resulting to a less sticky mucus. Ambroxol makes the mucus/ phlegm easier to cough out and breathing is eased considerably.

### Strength of the Medicine

Each sustained released capsule contains:

Ambroxol hydrochloride.....75mg

### What is this Medicine used for?

This medicine is used for the relief of cough caused by acute and chronic disorder of the respiratory tract accompanied with thick and sticky mucus such as acute and chronic bronchitis, asthmatoïd bronchitis and bronchial asthma.

### How much and how often should you use this Medicine?

Adult and Children over 12 years should take 1 capsule once daily after meal with plenty of liquid (sufficient supply of liquid supports the expectorant effect of Zobrixol). Or as prescribed by the physician.

### When should you not take this Medicine?

If you are allergic to any component in the product.

### Care that should be taken when taking this Medicine?

Do not take more than the recommended dose

Consult a physician if any adverse effect occurs.

Not recommended during the first three month of pregnancy.

Do not use after expiry date.

### Undesirable Effects of this Medicine

Ambroxol is generally well tolerated. Gastro-intestinal disturbances (diarrhea, nausea, vomiting), transient rise in serum aminotransferase. In rare cases weakness, headache and skin rash.

### What other Medicine or food should be avoided while taking this Medicine?

Ambroxol HCl (Zobrixol) should not be taken simultaneously with antitussive (e.g codeine), because it could lead to non-expectoration of phlegm or mucus previously liquified by Zobrixol.

### What should you do if you miss a dose?

If you miss a dose, just take the dose if still needed and subsequent doses at the usual recommended schedule. Do not exceed recommended dose.

### Signs and Symptoms of Overdose

Symptoms of Ambroxol overdose include extension of Undesirable effects such as Gastro-intestinal disturbances (diarrhea, nausea, vomiting) and skin rashes.

### What to do when you have taken more than the recommended dosage?

If you have taken more than the recommended dosage, consult a physician.

### How should you keep this Medicine?

Store at temperatures not exceeding 30°C. Keep out of reach of children.

### When should you consult your doctor?

If symptoms persist and if any undesirable effects occur.

### Availability

Aluminum Blister foil x 10's (Box of 50's)

### Manufactured for

Natrapharm Inc.

The Patriot Building Km 18, West Service Road SLEX, Sucat, Parañaque City.

### Name and Address of Manufacturer

Llody Laboratories Inc.

No. 10 Lloyd Avenue First Bulacan Industrial City, City of Malolos, Bulacan.

### ADR Reporting Statement

"For suspected adverse drug reaction, report to the FDA: [www.fda.gov.ph](http://www.fda.gov.ph)"

### Registration Number

DR-XY32403

### Date of First Authorization

October 2006

### Date of Revision of Patient Information Leaflet

June 2021

NTZBX0750001N2101